

On higher-power moments of $\Delta(x)$ (II)

Wenguang Zhai

School of Mathematical Sciences
Shandong Normal University
Jinan 250014, Shandong
P. R. China
zhaiwg@hotmail.com

Acta Arith. Vol.**114**(2004), 35-54

Abstract

Let $\Delta(x)$ be the error term of the Dirichlet divisor problem. The asymptotic formula of the integral $\int_1^T \Delta^k(x)dx$ is established for any integer $3 \leq k \leq 9$ by an unified method. Similar results are also established for some other well-known error terms in the analytic number theory .

1 Introduction and main results

1.1 Notations

Throughout this paper, let $d(n)$ denote the Dirichlet divisor function, $r(n)$ denote the number of ways n can be written as $n = x^2 + y^2$ for $x, y \in \mathbb{Z}$, and $a(n)$ denote the Fourier coefficients of a holomorphic cusp form of weight $\kappa = 2n \geq 12$ for the full modular group , $\tilde{a}(n) := a(n)n^{-\kappa/2+1/2}$. For short, we use d, r, a, \tilde{a} denote these functions, respectively. $\zeta(s)$ denotes the Riemann zeta-function.

2000 Mathematics Subject Classification: 11N37, 11M06.

Suppose $x > 0, t > 0$. Define

$$\Delta(x) := \sum_{n \leq x} d(n) - x \log x - (2\gamma - 1)x, \quad (1.1)$$

$$P(x) := \sum_{n \leq x} r(n) - \pi x, \quad (1.2)$$

$$A(x) := \sum_{n \leq x} a(n), \quad (1.3)$$

$$E(t) := \int_0^t |\zeta(\frac{1}{2} + iu)|^2 du - t \log(t/2\pi) - (2\gamma - 1)t. \quad (1.4)$$

Suppose $f : \mathbb{N} \rightarrow \mathbb{R}$ is any function such that $f(n) \ll n^\varepsilon$, $k \geq 2$ is a fixed integer. Define

$$s_{k;l}(f) := \sum_{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k}} \frac{f(n_1) \cdots f(n_k)}{(n_1 \cdots n_k)^{3/4}} \quad (1 \leq l < k), \quad (1.5)$$

$$B_k(f) := \sum_{l=1}^{k-1} \binom{k-1}{l} s_{k;l}(f) \cos \frac{\pi(k-2l)}{4}. \quad (1.6)$$

We shall use $s_{k;l}(f)$ to denote both of the series (1.5) and its value. We will prove the convergence of $s_{k;l}(f)$ in Section 3.

Suppose $A_0 > 2$ is a real number, define

$$\begin{aligned} K_0 &:= \min\{n \in \mathbb{N} : n \geq A_0, 2|n\}, \\ b(k) &:= 2^{k-2} + \frac{k-6}{4}, \\ \sigma(k, A_0) &:= \begin{cases} 1/4, & \text{if } k-1 < A_0/2, \\ \frac{A_0-k}{2(A_0-2)}, & \text{if } A_0/2 + 1 \leq k < A_0, \end{cases} \\ \delta_1(k, A_0) &:= \sigma(k, A_0)/2b(K_0), \\ \delta_2(k, A_0) &:= \frac{\sigma(k, A_0)}{2b(k) + 2\sigma(k, A_0)}. \end{aligned}$$

\mathbb{N} denotes the set of all natural numbers. ε always denotes a sufficiently small positive constant which may be different at different places. We will use the inequality $d(n) \ll n^\varepsilon$ freely. $SC(\Sigma)$ denotes the summation condition of the sum Σ . $\mu(n)$ is the Möbius function.

1.2 Introduction

In this paper we shall study the higher-power moments of $\Delta(x)$, $P(x)$, $A(x)$ and $E(t)$.

We begin with the Dirichlet divisor problem. Dirichlet first proved that $\Delta(x) = O(x^{1/2})$. The exponent $1/2$ was improved by many authors. The latest result reads

$$\Delta(x) \ll x^{23/73}(\log x)^{315/146}, \quad (1.7)$$

which can be found in Huxley[6]. It is conjectured that

$$\Delta(x) = O(x^{1/4+\varepsilon}), \quad (1.8)$$

which is supported by the classical mean-square result

$$\int_1^T \Delta^2(x) dx = \frac{(\zeta(3/2))^4}{6\pi^2\zeta(3)} T^{3/2} + O(T \log^5 T) \quad (1.9)$$

proved by Tong[17] and the upper bound estimate

$$\int_1^T |\Delta(x)|^{A_0} dx \ll T^{1+A_0/4+\varepsilon}, \quad (1.10)$$

where $A_0 > 2$ is a fixed real number. The estimate of type (1.10) can be found in Ivić[7, Thm. 13.9] with $A_0 = 35/4$ and Heath-Brown[5] with $A_0 = 28/3$. On the other hand, Voronoi[19] proved that

$$\int_1^T \Delta(x) dx = T/4 + O(T^{3/4}), \quad (1.11)$$

which in conjunction with (1.9) shows that $\Delta(x)$ has a lot of sign change s and cancelations between the positive and negative portions.

Tsang[18] first studied the third- and fourth-power moments of $\Delta(x)$. He proved that (with notations in Section 1.1)

$$\int_1^T \Delta^3(x) dx = \frac{3s_{3;1}(d)}{28\pi^3} T^{7/4} + O(T^{7/4-1/14+\varepsilon}), \quad (1.12)$$

$$\int_1^T \Delta^4(x) dx = \frac{3s_{4;2}(d)}{64\pi^4} T^2 + O(T^{2-1/23+\varepsilon}). \quad (1.13)$$

Heath-Brown[5] proved that for $k = 3, 4, 5, 6, 7, 8, 9$ the limit

$$\lim_{T \rightarrow \infty} T^{-1-k/4} \int_1^T \Delta(x)^k dx$$

exists.

In [20] the author improved Tsang's method and proved that

$$\int_1^T \Delta^3(x)dx = \frac{3s_{3;1}(d)}{28\pi^3}T^{7/4} + O(T^{3/2+\varepsilon}), \quad (1.14)$$

$$\int_1^T \Delta^4(x)dx = \frac{3s_{4;2}(d)}{64\pi^4}T^2 + O(T^{2-2/41}), \quad (1.15)$$

$$\int_1^T \Delta^5(x)dx = \frac{5(2s_{5;2}(d) - s_{5;1}(d))}{288\pi^5}T^{9/4} + O(T^{9/4-5/816}). \quad (1.16)$$

But the argument of [20] fails for $k \geq 6$.

1.3 New results on higher-power moments of $\Delta(x)$

In this paper we shall use a different approach to study the higher-power moments of $\Delta(x)$. This leads to the asymptotic formulas of the integral $\int_1^T \Delta^k(x)dx$ for $3 \leq k \leq 9$. Furthermore, if the estimate (1.8) is true, then our approach can give the asymptotic formulas of $\int_1^T \Delta^k(x)dx$ for any $k \geq 10$.

Theorem 1. Let $A_0 > 9$ be a real number such that (1.10) holds, then for any integer $3 \leq k < A_0$, we have the asymptotic formula

$$\int_1^T \Delta^k(x)dx = \frac{B_k(d)}{(1+k/4)2^{3k/2-1}\pi^k}T^{1+k/4} + O(T^{1+k/4-\delta_1(k,A_0)+\varepsilon}) \quad (1.17)$$

Remark 1.1. From Ivić's argument[7, Thm 13.9], we know that the value of A_0 for which (1.10) holds depends on the large-value estimate and the upper bound estimate of $\Delta(x)$. If we insert the estimate (1.7) into the argument of Ivić, we get that (1.10) holds with $A_0 = 184/19$. Whence for $k \in \{3, 4, 5, 6, 7, 8, 9\}$, we can get the asymptotic formula (1.17). Moreover, if the estimate $\Delta(x) \ll x^{5/16-\delta}$ holds for some small $\delta > 0$, then the asymptotic formula (1.17) holds for $k = 10$.

Remark 1.2. For $k \geq 10$, Theorem 1 is only an conditional result. However, it tells us that for any $k \geq 10$, the main term in the asymptotic formula of $\int_1^T \Delta^k(x)dx$ (if it exists) must have the form stated in (1.17).

Remark 1.3. We can state the following three conjectures about $\Delta(x)$.

Conjecture 1: The estimate (1.8) is true.

Conjecture 2: The estimate (1.10) is true for any $A_0 > 2$.

Conjecture 3: For any fixed $k \geq 3$, there exists a constant $\delta_k > 0$ such that the asymptotic formula

$$\int_1^T \Delta^k(x) dx = \frac{B_k(d)}{(1 + k/4)2^{3k/2-1}\pi^k} T^{1+k/4} + O(T^{1+k/4-\delta_k+\varepsilon})$$

holds.

It is well-known that Conjecture 1 and Conjecture 2 are equivalent. From Theorem 1 we know that actually the three Conjectures are equivalent. It is easy to deduce Conjecture 2 from Conjecture 3. To deduce Conjecture 3 from Conjecture 2, we take $A_0 = 2(k-1)$ and $\delta_k = \delta_1(k, 2(k-1))$.

Remark 1.4. From (1.11) we know that the integral $\int_1^T \Delta(x) dx$ have many cancelations from the positive and negative portions of $\Delta(x)$. However, from (1.12) Tsang[18] observed that this is not so for $\int_1^T \Delta^3(x) dx$. From Theorem 1 we know this is also not so for $\int_1^T \Delta^k(x) dx$ ($k = 5, 7, 9$) since numerical computation tells $B_k(d) > 0$ for $k = 5, 7, 9$. Maybe $B_k(d) > 0$ holds for any odd $k \geq 3$.

The constant $\delta_1(k, A_0)$ is small for k small. If we combine Ivić's argument in the proof of Theorem 1, we can get the following Theorem 2 for $3 \leq k \leq 9$. Note that the results for $k = 3, 4$ are weaker than those of [20]. Theorem 2 for $k = 5$ improved (1.16).

Theorem 2. For $3 \leq k \leq 9$, the asymptotic formula (1.17) holds with $\delta_1(k, A_0)$ replaced by $\delta_2(k, 184/19)$.

Especially for $k = 5, 6, 7, 8, 9$, we have

$$\int_1^T \Delta^5(x) dx = \frac{5(2s_{5;2}(d) - s_{5;1}(d))}{288\pi^5} T^{9/4} + O(T^{9/4-1/64+\varepsilon}), \quad (1.18)$$

$$\int_1^T \Delta^6(x) dx = \frac{5s_{6;3}(d) - 3s_{6;1}(d)}{320\pi^6} T^{5/2} + O(T^{5/2-35/4742+\varepsilon}), \quad (1.19)$$

$$\int_1^T \Delta^7(x) dx = \frac{7(5s_{7;3}(d) - 3s_{7;2}(d) - s_{7;1}(d))}{2816\pi^7} T^{11/4} + O(T^{11/4-17/6312+\varepsilon}), \quad (1.20)$$

$$\int_1^T \Delta^8(x) dx = \frac{7(5s_{8;4}(d) - 4s_{8;2}(d))}{6144\pi^8} T^3 + O(T^{3-8/9433+\varepsilon}), \quad (1.21)$$

$$\int_1^T \Delta^9(x) dx = \frac{3(3s_{9;1}(d) - 12s_{9;2}(d) - 28s_{9;3}(d) + 42s_{9;4}(d))}{26624\pi^9} T^{13/4} \quad (1.22)$$

$$+ O(T^{13/4-13/75216+\varepsilon}).$$

1.4 Higher-power moments of $P(x)$, $A(x)$ and $E(t)$

The method of proving Theorem 1 and Theorem 2 can also be applied to study the higher-power moments of $P(x)$, $A(x)$ and $E(t)$.

The conjectured bound of $P(x)$ is

$$P(x) = O(x^{1/4+\varepsilon}), \quad (1.23)$$

which is supported by

$$\int_2^T P^2(x)dx = \left(\frac{1}{3\pi^2} \sum_{n=1}^{\infty} r^2(n)n^{-3/2}\right)T^{3/2} + O(T \log^2 T) \quad (1.24)$$

proved by Katai[14]. Tsang[18] also studied the third- and the fourth-power moments of $P(x)$. His results were improved in the author[20]. An asymptotic formula for the fifth-power moment of $P(x)$ was also obtained in [20], which is further improved by the following Theorem 3($k = 5$.)

Theorem 3. Let $A_0 > 9$ be a real number such that the estimate

$$\int_1^T |P(x)|^{A_0} dx \ll T^{1+A_0/4+\varepsilon} \quad (1.25)$$

is true, then for any integer $3 \leq k < A_0$, the asymptotic formula

$$\int_1^T P^k(x)dx = \frac{(-1)^k B_k(r)}{(1+k/4)2^{k-1}\pi^k} T^{1+k/4} + O(T^{1+k/4-\delta_1(k,A_0)+\varepsilon}) \quad (1.26)$$

holds.

Especially for $3 \leq k \leq 9$, the asymptotic formula (1.26) holds with $\delta_1(k, A_0)$ replaced by $\delta_2(k, 184/19)$.

Remark 1.5. Ivić [7, Thm 13.12] proved that the estimate (1.25) holds for $A_0 = 35/4$. If we insert the estimate $P(x) = O(x^{23/73+\varepsilon})$ (see Huxley[6]) into his argument, we find that (1.25) holds for $A_0 = 184/19$.

It is well-known that $A(x)$ has no main term and $A(x) \ll x^{\kappa/2-1/6+\varepsilon}$. From Deligne[4], we have $|\tilde{a}(n)| \leq d(n)$.

The conjectured bound of $A(x)$ is $A(x) \ll x^{\kappa/2-1/4+\varepsilon}$. Ivić[9] proved that

$$\int_1^T A^2(x)dx = B_2 T^{\kappa+1/2} + O(T^{\kappa} \log^5 T), \quad (1.27)$$

where

$$B_2 = \frac{1}{4\kappa + 2} \sum_{n=1}^{\infty} a^2(n) n^{-\kappa-1/2}.$$

Ivić[9] also proved that the estimate

$$\int_1^T |A(x)|^{A_0} dx \ll T^{1+A_0(2\kappa-1)/4+\varepsilon} \quad (1.28)$$

holds for $A_0 = 8$. Cai [3] studied the third- and fourth-power moments of $A(x)$. His results were improved in the author[20]. In [20] an asymptotic formula for the fifth-power moment of $A(x)$ was also obtained, which is further improved by the case $k = 5$ of the following Theorem 4.

Theorem 4. Let $A_0 \geq 8$ be a real number such that (1.28) is true, then for any $3 \leq k < A_0$, the asymptotic formula

$$\int_1^T A^k(x) dx = \frac{B_k(\tilde{a})}{(1 + \frac{k(2\kappa-1)}{4}) 2^{3k/2-1} \pi^k} T^{1+\frac{k(2\kappa-1)}{4}} + O(T^{1+\frac{k(2\kappa-1)}{4}-\delta_1(k, A_0)+\varepsilon}) \quad (1.29)$$

holds.

Especially for $3 \leq k \leq 7$, the asymptotic formula (1.29) holds with $\delta_1(k, A_0)$ replaced by $\delta_2(k, 8)$.

Many results for $E(t)$ parallel to those for $\Delta(x)$ have been obtained; see Ivić[8] for a survey. The conjectured bound for $E(t)$ is $E(t) \ll t^{1/4+\varepsilon}$, which is supported by

$$\int_2^T E^2(t) dt = \frac{2\zeta^4(3/2)}{3\zeta(3)\sqrt{2\pi}} T^{3/2} + O(T \log^5 T), \quad (1.30)$$

proved by Meurman[15]. It has been proved that (see Huxley[6])

$$E(t) \ll t^{72/227} (\log t)^{629/227}, \quad t > 2. \quad (1.31)$$

Ivić[7, Thm. 15.7] proved that the estimate

$$\int_1^T |E(t)|^{A_0} dt \ll T^{1+A_0/4+\varepsilon} \quad (1.32)$$

holds for $A_0 = 35/4$. Inserting the estimate (1.31) into Ivić's argument, we find that (1.32) is true for $A_0 = 576/61$.

Tsang[18] studied the third- and fourth-power moment of $E(t)$ by using the Atkinson's formula[1]. His results were further improved by Ivić[10] in

a different way. The author[20] obtained new results on the third and the fourth power moments of $E(t)$. An asymptotic formula for the fifth power moment of $E(t)$ was also obtained in [20], which is further improved by the case $k = 5$ of the following Theorem 5..

Theorem 5. Let $A_0 > 9$ be a real number such that the estimates (1.10) and (1.32) hold, then for any $3 \leq k < A_0$, we have the asymptotic formula

$$\int_1^T E^k(t)dt = \frac{B_k(d)}{(1 + k/4)2^{3k/4-1}\pi^{k/4}} T^{1+k/4} + O(T^{1+k/4-\delta_1(k,A_0)+\varepsilon}). \quad (1.33)$$

Especially for $3 \leq k \leq 9$, the asymptotic formula (1.33) holds with $\delta_1(k, A_0)$ replaced by $\delta_2(k, 576/61)$.

Acknowledgement. The author deeply thanks the referee for his warm and valuable comments. The author also thanks Prof. Isao Wakabayashi for helpful discussions, through which the proof of Lemma 3.1 was further improved.

2 Some Preliminary Lemmas

We need the following Lemmas.

Lemma 2.1. The square-roots of squarefree numbers are linearly independent over the integers.

Proof. This is a classical result of Besicovitch[2]. □

Lemma 2.2. Suppose $k \geq 3$, $(i_1, \dots, i_{k-1}) \in \{0, 1\}^{k-1}$ such that

$$\sqrt{n_1} + (-1)^{i_1} \sqrt{n_2} + (-1)^{i_2} \sqrt{n_3} + \dots + (-1)^{i_{k-1}} \sqrt{n_k} \neq 0.$$

Then we have

$$|\sqrt{n_1} + (-1)^{i_1} \sqrt{n_2} + (-1)^{i_2} \sqrt{n_3} + \dots + (-1)^{i_{k-1}} \sqrt{n_k}| \gg \max(n_1, \dots, n_k)^{-(2^{k-2}-2^{-1})}.$$

Proof. The cases $k = 3, 4$ are Lemma 1 and Lemma 2 of Tsang[18], respectively. The proof for the general case is the same as the proof of Lemma 1 of [18]. We note that Heath-Brown[5] stated a similar result for k even. □

Lemma 2.3. Suppose $A, B \in \mathbb{R}$, $A \neq 0$, then

$$\int_T^{2T} \cos(A\sqrt{t} + B)dt \ll T^{1/2}|A|^{-1}.$$

Lemma 2.4. Suppose $k \geq 3$, $(i_1, \dots, i_{k-1}) \in \{0, 1\}^{k-1}$, $(i_1, \dots, i_{k-1}) \neq (0, \dots, 0)$, $1 < N_1, N_2, \dots, N_k$, $0 < \Delta \ll E^{1/2}$, $E = \max(N_1, N_2, \dots, N_k)$. Let

$$\mathcal{A} = \mathcal{A}(N_1, N_2, \dots, N_k; i_1, \dots, i_{k-1}; \Delta)$$

denote the number of solutions of the inequality

$$|\sqrt{n_1} + (-1)^{i_1}\sqrt{n_2} + (-1)^{i_2}\sqrt{n_3} + \dots + (-1)^{i_{k-1}}\sqrt{n_k}| < \Delta \quad (2.1)$$

with $N_j < n_j \leq 2N_j$, $1 \leq j \leq k$. Then

$$\mathcal{A} \ll \Delta E^{-1/2} N_1 N_2 \dots N_k + E^{-1} N_1 N_2 \dots N_k.$$

Proof. Without loss of generality, suppose $E = N_k$. If (n_1, \dots, n_k) satisfies (2.1), then

$$\sqrt{n_1} + (-1)^{i_1}\sqrt{n_2} + (-1)^{i_2}\sqrt{n_3} + \dots + (-1)^{i_{k-2}}\sqrt{n_{k-1}} = (-1)^{i_{k-1}+1}\sqrt{n_k} + \theta\Delta$$

for some $|\theta| < 1$. Whence we get

$$(\sqrt{n_1} + (-1)^{i_1}\sqrt{n_2} + (-1)^{i_2}\sqrt{n_3} + \dots + (-1)^{i_{k-2}}\sqrt{n_{k-1}})^2 = n_k + O(\Delta N_k^{1/2}).$$

Hence for fixed (n_1, \dots, n_{k-1}) , the number of n_k is $\ll 1 + \Delta N_k^{1/2}$ and thus

$$\mathcal{A} \ll \Delta N_k^{1/2} N_1 N_2 \dots N_{k-1} + N_1 N_2 \dots N_{k-1}.$$

□

3 On the series $s_{k;l}(d)$

In this section we shall study the series $s_{k;l}(d)$. Suppose $y > 1$ is a large parameter, and define

$$s_{k;l}(d; y) := \sum_{\substack{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k} \\ n_1, \dots, n_k \leq y}} \frac{d(n_1) \dots d(n_k)}{(n_1 \dots n_k)^{3/4}}, 1 \leq l < k.$$

We shall prove the following Lemma 3.1.

Lemma 3.1. We have

$$|s_{k;l}(d) - s_{k;l}(d; y)| \ll y^{-1/2+\varepsilon}, 1 \leq l < k.$$

Remark. Lemma 3.1 is still true if the divisor function d is replaced by any function $f : \mathbb{N} \rightarrow \mathbb{R}$ with $f(n) \ll n^\varepsilon$.

Proof. We shall prove Lemma 3.1 by induction in k . The case $k = 2$ is easy. The case $k = 3$ is contained already in Page 70 of Tsang[18], later we always suppose $k \geq 4$. Since $s_{k;l}(d) = s_{k;k-l}(d)$, we suppose $l \leq k/2$.

By the symmetry, we get

$$\begin{aligned} |s_{k;l}(d) - s_{k;l}(d; y)| &\ll \sum_{\substack{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k} \\ n_1 > y}} \frac{d(n_1) \cdots d(n_k)}{(n_1 \cdots n_k)^{3/4}} \\ &\ll U_1(d; y) + U_2(d; y), \end{aligned} \quad (3.1)$$

say, where

$$\begin{aligned} U_1(d; y) &:= \sum_{j=l+1}^k \sum_{\substack{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k} \\ n_1 = n_j > y}} \frac{d(n_1) \cdots d(n_k)}{(n_1 \cdots n_k)^{3/4}}, \\ U_2(d; y) &:= \sum_{\substack{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k} \\ n_1 > y, n_1 \neq n_j, l+1 \leq j \leq k}} \frac{d(n_1) \cdots d(n_k)}{(n_1 \cdots n_k)^{3/4}}. \end{aligned}$$

If $l = 1$, then obviously $U_1(d; y) = 0$. If $l > 1$, then by induction we get

$$U_1(d; y) \ll \sum_{n > y} \frac{d^2(n)}{n^{3/2}} s_{k-2;l-1}(d) \ll y^{-1/2+\varepsilon}. \quad (3.2)$$

Now we estimate $U_2(d; y)$. Let $I = \{1, \dots, l\}$, $J = \{l+1, \dots, k\}$. Suppose $(n_1, \dots, n_k) \in \mathbb{N}^k$ such that

$$(*) : \quad \sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k}, n_1 \neq n_j, l+1 \leq j \leq k.$$

Then there exist two sets $I_0 \subset I$, $J_0 \subset J$ which satisfy the following properties:

- (1). $1 \in I_0$;
- (2). $\sum_{i \in I_0} \sqrt{n_i} = \sum_{j \in J_0} \sqrt{n_j}$;
- (3). For any real subset $I'_0 \subset I_0$, $J'_0 \subset J_0$, we have

$$\sum_{i \in I'_0} \sqrt{n_i} \neq \sum_{j \in J'_0} \sqrt{n_j}.$$

If $(I_0, J_0) = (I, J)$, then we say (n_1, \dots, n_k) is a primitive (k, l) -point. Let $\mathbb{N}_{k;l}$ denote the set of all points in \mathbb{N}^k which satisfy $(*)$ and $\mathbb{N}_{k;l}^*$ the set of all

primitive (k, l) -points, respectively. Let $\mathcal{G}_{k;l}$ denote the set of all possible pairs (I_0, J_0) when (n_1, \dots, n_k) runs through $\mathbb{N}_{k;l}$. Note that if $l = 1$, then $\mathcal{G}_{k;l} = \{(I, J)\}$.

Suppose $(I_0, J_0) \in \mathcal{G}_{k;l}$. Let $l_1 = \#I_0, l_2 = l - l_1, k_1 = \#I_0 + \#J_0, k_2 = k - k_1$. From (*), we know that $k_1 \geq 3$. Define

$$R_1^{(I_0, J_0)}(d; y) := \sum_{\substack{\sqrt{n_1} + \dots + \sqrt{n_{l_1}} = \sqrt{n_{l_1+1}} + \dots + \sqrt{n_{k_1}} \\ n_1 > y, (n_1, \dots, n_{k_1}) \in \mathbb{N}_{k_1; l_1}^*}} \frac{d(n_1) \cdots d(n_{k_1})}{(n_1 \cdots n_{k_1})^{3/4}}.$$

If $(I_0, J_0) \neq (I, J)$, then $l_1 < l, k_1 < k$ and we define

$$R_2^{(I_0, J_0)}(d) := \sum_{\sqrt{m_1} + \dots + \sqrt{m_{l_2}} = \sqrt{m_{l_2+1}} + \dots + \sqrt{m_{k_2}}} \frac{d(m_1) \cdots d(m_{k_2})}{(m_1 \cdots m_{k_2})^{3/4}}.$$

By the induction assumption, $R_2^{(I_0, J_0)}(d) \ll 1$.

If $(n_1, \dots, n_{k_1}) \in \mathbb{N}_{k_1; l_1}^*$, then by Lemma 2.1 we have

$$n_j = s_j^2 h, s_1 + \dots + s_{l_1} = s_{l_1+1} + \dots + s_{k_1}, \mu(h) \neq 0.$$

$n_1 > y$ implies that there exists at least one $n_j (l_1 + 1 \leq j \leq j_1)$ such that $n_j \gg y$. We suppose $n_{k_1} \gg y$. So we have

$$\begin{aligned} R_1^{(I_0, J_0)}(d; y) &\ll \sum_h \sum_{\substack{s_1 + \dots + s_{l_1} = s_{l_1+1} + \dots + s_{k_1} \\ s_1^2 h > y, s_{k_1}^2 h \gg y}} \frac{d(s_1^2 h) \cdots d(s_{k_1}^2 h)}{h^{3k_1/4} (s_1 \cdots s_{k_1})^{3/2}} \\ &\ll \sum_h \sum_{\substack{s_1 + \dots + s_{l_1} = s_{l_1+1} + \dots + s_{k_1} \\ s_1^2 h > y, s_{k_1}^2 h \gg y}} \frac{d^2(s_1) \cdots d^2(s_{k_1}) d^{k_1}(h)}{h^{3k_1/4} (s_1 \cdots s_{k_1})^{3/2}} \\ &\ll \sum_h \frac{d^{k_1}(h)}{h^{3k_1/4}} \sum_{s_1 > (y/h)^{1/2}} \frac{d^2(s_1)}{s_1^{3/2}} \sum_{s_{k_1} \gg (y/h)^{1/2}} \frac{d^2(s_{k_1})}{s_{k_1}^{3/2}} \\ &\ll \sum_h \frac{d^{k_1}(h)}{h^{3k_1/4}} \left(\frac{y}{h}\right)^{-1/2+\varepsilon} \ll y^{-1/2+\varepsilon} \end{aligned}$$

if we notice $k_1 \geq 3$.

If $\mathcal{G}_{k;l} = (I, J)$, we have

$$U_2(d; y) \ll R_1^{(I, J)}(d; y) \ll y^{-1/2+\varepsilon}. \quad (3.3)$$

If $\mathcal{G}_{k;l} \neq (I, J)$, we have

$$U_2(d; y) \ll R_1^{(I, J)}(d; y) + \sum_{\substack{(I_0, J_0) \in \mathcal{G}_{k;l} \\ (I_0, J_0) \neq (I, J)}} R_1^{(I_0, J_0)}(d; y) R_2^{(I_0, J_0)}(d) \ll y^{-1/2+\varepsilon}. \quad (3.4)$$

Now Lemma 3.1 follows from (3.1)-(3.4). \square

4 Proofs of Theorem 1 and Theorem 2

Suppose $T \geq 10$ is a real number. It suffices for us to evaluate the integral $\int_T^{2T} \Delta^k(x) dx$. Suppose y is a parameter such that $T^\varepsilon < y \leq T^{1/3}$. For any $T \leq x \leq 2T$, define

$$\begin{aligned} \mathcal{R}_1 &= \mathcal{R}_1(x, y) := (\sqrt{2}\pi)^{-1} x^{1/4} \sum_{n \leq y} \frac{d(n)}{n^{3/4}} \cos(4\pi\sqrt{xn} - \frac{\pi}{4}), \\ \mathcal{R}_2 &= \mathcal{R}_2(x, y) := \Delta(x) - \mathcal{R}_1. \end{aligned}$$

We shall show that the higher-power moment of \mathcal{R}_2 is small and hence the integral $\int_T^{2T} \Delta^k(x) dx$ can be well approximated by $\int_T^{2T} \mathcal{R}_1^k dx$, which is easy to evaluate.

4.1 Evaluation of the integral $\int_T^{2T} \mathcal{R}_1^h dx$

Suppose $h \geq 3$ is any fixed integer. By the elementary formula

$$\cos a_1 \cdots \cos a_h = \frac{1}{2^{h-1}} \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \cos(a_1 + (-1)^{i_1} a_2 + (-1)^{i_2} a_3 + \cdots + (-1)^{i_{h-1}} a_h),$$

we have

$$\begin{aligned} \mathcal{R}_1^h &= (\sqrt{2}\pi)^{-h} x^{\frac{h}{4}} \sum_{n_1 \leq y} \cdots \sum_{n_h \leq y} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}} \prod_{j=1}^h \cos(4\pi\sqrt{n_j x} - \pi/4) \\ &= \frac{x^{\frac{h}{4}}}{(\sqrt{2}\pi)^h 2^{h-1}} \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \sum_{n_1 \leq y} \cdots \sum_{n_h \leq y} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}} \\ &\quad \times \cos(4\pi\sqrt{x}\alpha(n_1, \dots, n_h; i_1, \dots, i_{h-1}) - \frac{\pi}{4}\beta(i_1, \dots, i_{h-1})), \end{aligned}$$

where

$$\begin{aligned}\alpha(n_1, \dots, n_h; i_1, \dots, i_{h-1}) \\ &:= \sqrt{n_1} + (-1)^{i_1} \sqrt{n_2} + (-1)^{i_2} \sqrt{n_3} + \dots + (-1)^{i_{h-1}} \sqrt{n_h}, \\ \beta(i_1, \dots, i_{h-1}) &:= 1 + (-1)^{i_1} + (-1)^{i_2} + \dots + (-1)^{i_{h-1}}.\end{aligned}$$

Thus we can write

$$\mathcal{R}_1^h = \frac{1}{(\sqrt{2}\pi)^h 2^{h-1}} (S_1(x) + S_2(x)), \quad (4.1)$$

where

$$\begin{aligned}S_1(x) : &= x^{h/4} \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \cos\left(-\frac{\pi\beta}{4}\right) \sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha=0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}}, \\ S_2(x) : &= x^{h/4} \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha \neq 0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}} \cos\left(4\pi\alpha\sqrt{x} - \frac{\pi\beta}{4}\right), \\ \alpha : &= \alpha(n_1, \dots, n_h; i_1, \dots, i_{h-1}), \\ \beta : &= \beta(i_1, \dots, i_{h-1}).\end{aligned}$$

First consider the contribution of $S_1(x)$. We have

$$\int_T^{2T} S_1(x) dx = \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \cos\left(-\frac{\pi\beta}{4}\right) \sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha=0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}} \int_T^{2T} x^{\frac{h}{4}} dx. \quad (4.2)$$

It is easily seen that if $\alpha = 0$, then $1 \in \{i_1, \dots, i_{h-1}\}$. Let $l = i_1 + \dots + i_{h-1}$, then we have

$$\sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha=0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}} = s_{h,l}(d; y),$$

where $s_{h,l}(d; y)$ was defined in last section. By Lemma 3.1 we get

$$\int_T^{2T} S_1(x) dx = B_h^*(d) \int_T^{2T} x^{\frac{h}{4}} dx + O(T^{1+h/4+\varepsilon} y^{-1/2}), \quad (4.3)$$

where

$$B_h^*(d) := \sum_{(i_1, \dots, i_{h-1}) \in \{0,1\}^{h-1}} \cos\left(-\frac{\pi\beta}{4}\right) \sum_{\substack{(n_1, \dots, n_h) \in \mathbb{N}^h \\ \alpha=0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}}.$$

For any $(i_1, \dots, i_{h-1}) \in \{0, 1\}^{h-1} \setminus \{(0, \dots, 0)\}$, let

$$\begin{aligned} S(d; i_1, \dots, i_{h-1}) : &= \sum_{\substack{(n_1, \dots, n_h) \in \mathbb{N}^h \\ \alpha=0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4}}, \\ l(i_1, \dots, i_{h-1}) : &= i_1 + \cdots + i_{h-1}. \end{aligned}$$

It is easily seen that if $l(i_1, \dots, i_{h-1}) = l(i'_1, \dots, i'_{h-1})$ or $l(i_1, \dots, i_{h-1}) + l(i'_1, \dots, i'_{h-1}) = h$, then

$$S(d; i_1, \dots, i_{h-1}) = S(d; i'_1, \dots, i'_{h-1}) = s_{h; l(i_1, \dots, i_{h-1})}(d).$$

From $(-1)^i = 1 - 2i(i = 0, 1)$ we also have

$$\beta(i_1, \dots, i_{h-1}) = h - 2l(i_1, \dots, i_{h-1}).$$

So we get

$$\begin{aligned} B_h^*(d) &= \sum_{l=1}^{h-1} \sum_{l(i_1, \dots, i_{h-1})=l} \cos\left(-\frac{\pi\beta}{4}\right) S(d; i_1, \dots, i_{h-1}) \\ &= \sum_{l=1}^{h-1} s_{h;l}(d) \cos \frac{\pi(h-2l)}{4} \sum_{l(i_1, \dots, i_{h-1})=l} 1 \\ &= \sum_{l=1}^{h-1} \binom{h-1}{l} s_{h;l}(d) \cos \frac{\pi(h-2l)}{4} = B_h(d). \end{aligned} \tag{4.4}$$

Now we consider the contribution of $S_2(x)$. By Lemma 2.3 we get

$$\int_T^{2T} S_2(x) dx \ll T^{1/2+h/4} \sum_{(i_1, \dots, i_{h-1}) \in \{0, 1\}^{h-1}} \sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha \neq 0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4} |\alpha|}. \tag{4.5}$$

It suffices for us to estimate the sum

$$\Sigma(y; i_1, \dots, i_{h-1}) = \sum_{\substack{n_j \leq y, 1 \leq j \leq h \\ \alpha \neq 0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4} |\alpha|}$$

for fixed $(i_1, \dots, i_{h-1}) \in \{0, 1\}^{h-1}$.

If $(i_1, \dots, i_{h-1}) = (0, \dots, 0)$, then

$$\begin{aligned} \Sigma(y; 0, \dots, 0) &\ll \sum_{n_j \leq y, 1 \leq j \leq h} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4} (\sqrt{n_1} + \cdots + \sqrt{n_h})} \\ &\ll \sum_{n_j \leq y, 1 \leq j \leq h} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4 + 1/2h}} \\ &\ll y^{(h-2)/4} \log^h y, \end{aligned}$$

where we used the estimates

$$\sum_{n \leq u} \ll u \log u, \quad x_1 + \cdots + x_h \gg (x_1 \cdots x_h)^{1/h}.$$

For $(i_1, \dots, i_{h-1}) \neq (0, \dots, 0)$, by a splitting argument we get that there exist a group of numbers $1 < N_1, N_2, \dots, N_h < y$ such that

$$\Sigma(y; i_1, \dots, i_{h-1}) \ll \Sigma_1^* \log^h y,$$

where

$$\Sigma_1^* = \sum_{\substack{N_j < n_j \leq 2N_j, 1 \leq j \leq h \\ \alpha \neq 0}} \frac{d(n_1) \cdots d(n_h)}{(n_1 \cdots n_h)^{3/4} |\alpha|}.$$

Without loss of generality, we suppose $N_1 \leq N_2 \leq \cdots \leq N_h \leq y$.

By Lemma 2.2 we have $|\alpha| \gg N_h^{-(2^{h-2}-2^{-1})}$. Then by a splitting argument and Lemma 2.4 we get for some $N_h^{-(2^{h-2}-2^{-1})} \ll \Delta < y^{1/2}$ such that

$$\begin{aligned} \Sigma_1^* &\ll \frac{y^\varepsilon}{(N_1 \cdots N_h)^{3/4} \Delta} \mathcal{A}(N_1, \dots, N_h; i_1, \dots, i_{h-1}; \Delta) \\ &\ll \frac{y^\varepsilon}{(N_1 \cdots N_h)^{3/4} \Delta} (\Delta N_h^{1/2} N_1 \cdots N_{h-1} + N_1 \cdots N_{h-1}) \\ &\ll y^\varepsilon \left(\frac{(N_1 \cdots N_{h-1})^{1/4}}{N_h^{1/4}} + \frac{(N_1 \cdots N_{h-1})^{1/4}}{N_h^{3/4} \Delta} \right) \\ &\ll y^\varepsilon (N_h^{(h-2)/4} + N_h^{b(h)}) \ll y^{b(h)+\varepsilon}, \end{aligned}$$

where $b(h)$ was defined in Section 1.1. Thus we get

$$\int_T^{2T} S_2(x) dx \ll T^{1/2+h/4+\varepsilon} y^{b(h)}. \quad (4.6)$$

Hence from (4.1)-(4.6) we get

Lemma 4.1. For any fixed $h \geq 3$, we have

$$\int_T^{2T} \mathcal{R}_1^h dx = \frac{B_h(d)}{(\sqrt{2}\pi)^h 2^{h-1}} \int_T^{2T} x^{h/4} dx + O(T^{1+h/4+\varepsilon} y^{-1/2} + T^{1/2+h/4+\varepsilon} y^{b(h)}). \quad (4.7)$$

4.2 Higher-power moments of \mathcal{R}_2

We first study the mean-square of \mathcal{R}_2 . We begin with the truncated Voronoi's formula[9, equation (2.25)]

$$\Delta(x) = (\pi\sqrt{2})^{-1} x^{\frac{1}{4}} \sum_{n \leq N} \frac{d(n)}{n^{3/4}} \cos(4\pi\sqrt{nx} - \pi/4) + O(x^{1/2+\varepsilon} N^{-1/2}), \quad (4.8)$$

where $1 < N \ll x$. Taking $N = T$, we get

$$\begin{aligned} \mathcal{R}_2 &= (\pi\sqrt{2})^{-1} x^{\frac{1}{4}} \sum_{y < n \leq T} \frac{d(n)}{n^{3/4}} \cos(4\pi\sqrt{nx} - \pi/4) + O(T^\varepsilon) \\ &\ll |x^{\frac{1}{4}} \sum_{y < n \leq T} \frac{d(n)}{n^{3/4}} e(2\sqrt{nx})| + T^\varepsilon, \end{aligned}$$

which implies

$$\begin{aligned} \int_T^{2T} \mathcal{R}_2^2 dx &\ll T^{1+\varepsilon} + \int_T^{2T} |x^{\frac{1}{4}} \sum_{y < n \leq T} \frac{d(n)}{n^{3/4}} e(2\sqrt{nx})|^2 dx \\ &\ll T^{1+\varepsilon} + T^{3/2} \sum_{y < n \leq T} \frac{d^2(n)}{n^{3/2}} + T \sum_{y < m < n \leq T} \frac{d(n)d(m)}{(mn)^{3/4}(\sqrt{n} - \sqrt{m})} \\ &\ll T^{1+\varepsilon} + \frac{T^{3/2} \log^3 T}{y^{1/2}} \ll \frac{T^{3/2} \log^3 T}{y^{1/2}}, \end{aligned} \quad (4.9)$$

where we used the estimates

$$\begin{aligned} \sum_{n \leq u} d^2(n) &\ll u \log^3 u, \\ \sum_{y < m < n \leq T} \frac{d(n)d(m)}{(mn)^{3/4}(\sqrt{n} - \sqrt{m})} &\ll T^\varepsilon. \end{aligned}$$

Now suppose y satisfies $y^{2b(K_0)} \leq T$. Hence from Lemma 4.1 we get that

$$\int_T^{2T} |\mathcal{R}_1|^{K_0} dx \ll T^{1+K_0/4+\varepsilon},$$

which implies

$$\int_T^{2T} |\mathcal{R}_1|^{A_0} dx \ll T^{1+A_0/4+\varepsilon} \quad (4.10)$$

since $A_0 \leq K_0$. From (1.10) and (4.10) we get

$$\int_T^{2T} |\mathcal{R}_2|^{A_0} dx \ll \int_T^{2T} (|\Delta(x)|^{A_0} + |\mathcal{R}_1|^{A_0}) dx \ll T^{1+A_0/4+\varepsilon}. \quad (4.11)$$

For any $2 < A < A_0$, from (4.9), (4.11) and Hölder's inequality we get that

$$\begin{aligned} \int_T^{2T} |\mathcal{R}_2|^A dx &= \int_T^{2T} |\mathcal{R}_2|^{\frac{2(A_0-A)}{A_0-2} + \frac{A_0(A-2)}{A_0-2}} dx \\ &\ll \left(\int_T^{2T} \mathcal{R}_2^2 dx \right)^{\frac{A_0-A}{A_0-2}} \left(\int_T^{2T} |\mathcal{R}_2|^{A_0} dx \right)^{\frac{A-2}{A_0-2}} \ll T^{1+\frac{A}{4}+\varepsilon} y^{-\frac{A_0-A}{2(A_0-2)}}. \end{aligned} \quad (4.12)$$

Namely, we have the following Lemma 4.2.

Lemma 4.2. Suppose $T^\varepsilon \leq y \leq T^{1/2b(K_0)}$, $2 < A < A_0$, then

$$\int_T^{2T} |\mathcal{R}_2|^A dx \ll T^{1+\frac{A}{4}+\varepsilon} y^{-\frac{A_0-A}{2(A_0-2)}}. \quad (4.13)$$

4.3 Proof of Theorem 1.

Suppose $3 \leq k \leq K(A_0)$ and $T^\varepsilon \leq y \leq T^{1/2b(K_0)}$. By the elementary formula $(a+b)^k - a^k \ll |a^{k-1}b| + |b|^k$, we get

$$\int_T^{2T} \Delta^k(x) dx = \int_T^{2T} \mathcal{R}_1^k dx + O\left(\int_T^{2T} |\mathcal{R}_1^{k-1} \mathcal{R}_2| dx\right) + O\left(\int_T^{2T} |\mathcal{R}_2|^k dx\right). \quad (4.14)$$

If $k-1 < A_0/2$, then from (4.9), (4.10) and Cauchy's inequality we get

$$\int_T^{2T} |\mathcal{R}_1^{k-1} \mathcal{R}_2| dx \ll \left(\int_T^{2T} |\mathcal{R}_1|^{2(k-1)} dx \right)^{1/2} \left(\int_T^{2T} |\mathcal{R}_2|^2 dx \right)^{1/2} \ll T^{1+k/4+\varepsilon} y^{-1/4}.$$

If $k - 1 \geq A_0/2$, then from (4.10), Lemma 4.2 and Hölder's inequality we get

$$\begin{aligned} \int_T^{2T} |\mathcal{R}_1^{k-1} \mathcal{R}_2| dx &\ll \left(\int_T^{2T} |\mathcal{R}_1|^{A_0} dx \right)^{\frac{k-1}{A_0}} \left(\int_T^{2T} |\mathcal{R}_2|^{\frac{A_0}{A_0-k+1}} dx \right)^{\frac{A_0-k+1}{A_0}} \\ &\ll T^{1+k/4+\varepsilon} y^{-(A_0-k)/2(A_0-2)}. \end{aligned}$$

Thus we have

$$\int_T^{2T} |\mathcal{R}_1^{k-1} \mathcal{R}_2| dx + \int_T^{2T} |\mathcal{R}_2|^k dx \ll T^{1+k/4+\varepsilon} y^{-\sigma(k, A_0)}, \quad (4.15)$$

where $\sigma(k, A_0)$ was defined in Section 1.1.

From (4.14) and (4.15) we get

$$\int_T^{2T} \Delta^k(x) dx = \int_T^{2T} \mathcal{R}_1^k dx + O(T^{1+k/4+\varepsilon} y^{-\sigma(k, A_0)}). \quad (4.16)$$

Now take $y = T^{1/2b(K_0)}$. From Lemma 4.1 and (4.16) we get

$$\begin{aligned} \int_T^{2T} \Delta^k(x) dx &= \frac{B_k(d)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{k/4} dx + O(T^{1+k/4-\sigma(k, A_0)/2b(K_0)+\varepsilon}) \\ &= \frac{B_k(d)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{k/4} dx + O(T^{1+k/4-\delta_1(k, A_0)+\varepsilon}). \end{aligned} \quad (4.17)$$

Theorem 1 follows from (4.17) immediately.

4.4 Proof of Theorem 2

Suppose $T^\varepsilon \leq y \leq T^{1/3}$. By the truncated Voronoi's formula (4.8), we have

$$\mathcal{R}_2 = (\sqrt{2}\pi)^{-1} x^{1/4} \sum_{y < n \leq N} \frac{d(n)}{n^{3/4}} \cos(4\pi\sqrt{nx} - \pi/4) + O(x^{1/2+\varepsilon} N^{-1/2}),$$

where $y < N \ll T$. Using Ivić's large-value technique directly to \mathcal{R}_2 without modifications, we get that the estimate

$$\int_T^{2T} |\mathcal{R}_2|^{A_0} dx \ll T^{1+A_0/4+\varepsilon} \quad (4.18)$$

holds with $A_0 = 184/19, T^\varepsilon \leq y \leq T^{1/3}$. We omit the details since it is completely the same as that of Ivić. Combining (1.10) we get that

$$\int_T^{2T} |\mathcal{R}_1|^{A_0} dx \ll T^{1+A_0/4+\varepsilon} \quad (4.19)$$

holds with $A_0 = 184/19, T^\varepsilon \leq y \leq T^{1/3}$.

By the same argument as in last subsection, we get that for $T^\varepsilon \leq y \leq T^{1/3}$,

$$\int_T^{2T} \Delta^k(x) dx = \int_T^{2T} \mathcal{R}_1^k dx + O(T^{1+k/4+\varepsilon} y^{-\sigma(k, 184/19)}). \quad (4.20)$$

Take $y = T^{1/(2b(k)+2\sigma(k, 184/19))}$. From Lemma 4.1 again we get

$$\begin{aligned} \int_T^{2T} \Delta^k(x) dx &= \frac{B_k(d)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{k/4} dx + O(T^{1+k/4-\frac{\sigma(k, 184/19)}{2b(k)+2\sigma(k, 184/19)}+\varepsilon}) \\ &= \frac{B_k(d)}{(\sqrt{2}\pi)^k 2^{k-1}} \int_T^{2T} x^{k/4} dx + O(T^{1+k/4-\delta_2(k, 184/19)+\varepsilon}). \end{aligned} \quad (4.21)$$

And Theorem 2 follows.

5 Proofs of other Theorems

$P(x)$ has the following truncated Voronoi's formula

$$P(x) = -\frac{1}{\pi} \sum_{n \leq N} r(n) n^{-3/4} x^{1/4} \cos(4\pi\sqrt{nx} + \pi/4) + O(x^{1/2+\varepsilon} N^{-1/2}) \quad (5.1)$$

for $1 \leq N \ll x$, which follows from Lemma 3 of Müller[16]. $A(x)$ has the following truncated Voronoi's formula

$$\begin{aligned} A(x) &= \frac{1}{\pi\sqrt{2}} x^{\kappa/2-1/4} \sum_{n \leq N} a(n) n^{-\kappa/2-1/4} \cos(4\pi\sqrt{nx} - \pi/4) \\ &\quad + O(x^{\kappa/2+\varepsilon} N^{-1/2}) \end{aligned} \quad (5.2)$$

for $1 \leq N \ll x$, which is a special case of Theorem 1.1 of Jutila[13]. So by the same way as in last section, we get Theorem 3 and Theorem 4.

Now we prove Theorem 5. We shall follow Ivić[10]. Define

$$\Delta^*(x) := \frac{1}{2} \sum_{n \leq 4x} (-1)^n d(n) - x(\log x + 2\gamma - 1), x > 0.$$

Jutila[12] proved that

$$\int_0^T (E(t) - 2\pi\Delta^*(\frac{t}{2\pi}))^2 dt \ll T^{4/3} \log^3 T, \quad (5.3)$$

which means that $E(t)$ is well approximated by $2\pi\Delta^*(\frac{t}{2\pi})$ at least in the mean square sense.

Suppose $A_0 > 9$ is a real number such that both of (1.10) and (1.32) hold. Since (see Jutila[11])

$$\Delta^*(x) = -\Delta(x) + 2\Delta(2x) - \frac{1}{2}\Delta(4x),$$

from (1.10) we get

$$\int_0^T |\Delta^*(t)|^{A_0} dt \ll T^{1+A_0/4+\varepsilon}. \quad (5.4)$$

Then from (1.32), (5.3), (5.4) and Hölder's inequality we get for any $3 \leq k < A_0$ that

$$\begin{aligned} & \int_0^T E^k(t) dt - (2\pi)^{k+1} \int_0^{\frac{T}{2\pi}} (\Delta^*(t))^k dt \\ &= \int_0^T \left(E^k(t) - (2\pi\Delta^*(\frac{t}{2\pi}))^k \right) dt \\ &\ll \int_0^T |E(t) - 2\pi\Delta^*(\frac{t}{2\pi})| \left(|E(t)|^{k-1} + |\Delta^*(\frac{t}{2\pi})|^{k-1} \right) dt \\ &\ll T^{1+k/4-\sigma(k, A_0)/3+\varepsilon}, \end{aligned} \quad (5.5)$$

where $\sigma(k, A_0)$ was defined in Section 1.1.

From (5.5) the problem is reduced to evaluating the integral $\int_0^T (\Delta^*(t))^k dt$. For $1 \ll N \ll x$, we have[10, equation (7)]

$$\Delta^*(x) = \frac{1}{\pi\sqrt{2}} \sum_{n \leq N} (-1)^n d(n) n^{-3/4} x^{1/4} \cos(4\pi\sqrt{nx} - \pi/4) + O(x^{1/2+\varepsilon} N^{-1/2}), \quad (5.6)$$

which is similar to (4.8). Let $d^*(n) = (-1)^n d(n)$. Then by the same way as in the proof of Theorem 1, we get that the asymptotic formula

$$\int_1^T (\Delta^*(t))^k dt = \frac{B_k(d^*)}{(1+k/4)2^{3k/2-1}\pi^k} T^{1+k/4} + O(T^{1+k/4-\delta_1(k, A_0)+\varepsilon}) \quad (5.7)$$

holds for any $3 \leq k < A_0$.

We shall use the following Lemma 5.1.

Lemma 5.1 Suppose $1 \leq l < k$ are fixed integers, $(n_1, \dots, n_k) \in \mathbb{N}^k$. If

$$\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k}$$

holds, then $2|(n_1 + \dots + n_k)$.

Proof. For any $n \in \mathbb{N}$, let $h(n)$ denote the squarefree part of n . Let $\mathcal{S} = \{h(n_1), \dots, h(n_k)\} \cap \mathbb{N}$ and $s = \#\mathcal{S}$. For convenience, write

$$\mathcal{S} = \{h_1, \dots, h_s\}, I = \{1, \dots, l\}, J = \{l+1, \dots, k\}.$$

From Lemma 2.1 we can write $I = \bigcup_{e=1}^s I_e, J = \bigcup_{e=1}^s J_e$ such that for each $1 \leq e \leq s$, we have

$$\sum_{i \in I_e} \sqrt{n_i} = \sum_{j \in J_e} \sqrt{n_j}$$

and that all $n_i (i \in I_e)$ and $n_j (j \in J_e)$ have the same squarefree part h_e . Namely we have ($1 \leq e \leq s$)

$$n_i = m_i^2 h_e (i \in I_e), n_j = m_j^2 h_e (j \in J_e), \sum_{i \in I_e} m_i = \sum_{j \in J_e} m_j.$$

Thus we get

$$\begin{aligned} n_1 + \dots + n_k &= \sum_{e=1}^s \left(\sum_{i \in I_e} n_i + \sum_{j \in J_e} n_j \right) \\ &= \sum_{e=1}^s \left(\sum_{i \in I_e} m_i^2 h_e + \sum_{j \in J_e} m_j^2 h_e \right) \equiv \sum_{e=1}^s \left(\sum_{i \in I_e} m_i + \sum_{j \in J_e} m_j \right) h_e \\ &= 2 \sum_{e=1}^s h_e \sum_{i \in I_e} m_i \equiv 0 \pmod{2}, \end{aligned}$$

where we used the simple congruence $n^2 \equiv n \pmod{2}$. □

From Lemma 5.1 we get for any $1 \leq l < k$ that

$$\begin{aligned} s_{k;l}(d^*) &= \sum_{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k}} (-1)^{n_1 + \dots + n_k} \frac{d(n_1) \dots d(n_k)}{(n_1 \dots n_k)^{3/4}} \\ &= \sum_{\sqrt{n_1} + \dots + \sqrt{n_l} = \sqrt{n_{l+1}} + \dots + \sqrt{n_k}} \frac{d(n_1) \dots d(n_k)}{(n_1 \dots n_k)^{3/4}} \\ &= s_{k;l}(d). \end{aligned}$$

Whence we get

$$B_k(d^*) = B_k(d). \quad (5.8)$$

From (5.5), (5.7) and (5.8) we get (1.33).

Similar to Theorem 2, we can prove that the asymptotic formula

$$\int_1^T (\Delta^*(t))^k dt = \frac{B_k(d)}{(1 + k/4)2^{3k/2-1}\pi^k} T^{1+k/4} + O(T^{1+k/4-\delta_2(k, 576/61)+\varepsilon}) \quad (5.9)$$

holds for any $3 \leq k \leq 9$, which combined with (5.5) yields the second part of Theorem 3.

Note added in proof: Recently M. N. Huxley (Exponential sums and Lattice points III, Proc. London Math. Soc., Vol.**87**(3)(2003), 591-609) proved

$$\Delta(x) \ll x^{131/416}(\log x)^{26947/8320},$$

which implies that the formula (1.10) holds for $A_0 = 262/27$. The exponent $\delta_2(k, 184/19)$ in Theorem 2 then can be improved to $\delta_2(k, 262/27)$ for $k = 6, 7, 8, 9$. The author deeply thanks Professor A. Schinzel for informing me M. N. Huxley's new result.

References

- [1] F. V. Atkinson, the mean value of the Riemann zeta-function, Acta Math.**81**(1949), 353-376.
- [2] A. S. Besicovitch, On the linear independence of fractional powers of integers, J. London Math. Soc.**15**(1940),3-6.
- [3] Cai Yingchun, On the third and fourth power moments of Fourier coefficients of cusp forms. Acta Math. Sinica (N.S.) 13 (1997), no. 4, 443–452.
- [4] P. Deligne, La conjecture de Weil I. Inst. Hautes Etudes Sci. Publ. Math.**43**(1974), 273-307.
- [5] D. R. Heath-Brown, The distribution and moments of the error term in the Dirichlet divisor problem, Acta Arith. (1992), 389-415.
- [6] M. N. Huxley, Area, lattice points, and exponential sums. London Math. Soc. Monographs. New Series, 13. Oxford University Press, 1996.
- [7] A. Ivić, The Riemann zeta-function. John. Wiley and Sons, 1985.

- [8] A. Ivić, Lectures on mean values of the Riemann zeta-function, Lectures On Math. and Physics 82, Tata Inst. Fund. Res., Bombay, 1991.
- [9] A. Ivić, Large values of certain number-theoretic error terms Acta Arith. **56**(1990), 135-159.
- [10] A. Ivić, On some problems involving the mean square of $\zeta(\frac{1}{2} + it)$. Bull. Cl. Sci. Math. Nat. Sci. Math. No. 23 (1998), 71–76.
- [11] M. Jutila, Riemann's zeta-function and the divisor problem Ark. Mat. **21**(1983), 75-96 and II, ibid. **31**(1993), 61-70.
- [12] M. Jutila, On a formula of Atkinson. Topics in classical number theory, Vol. I, II (Budapest, 1981), 807–823, Colloq. Math. Soc. Janos Bolyai, 34, North-Holland, Amsterdam, 1984.
- [13] M. Jutila, Lectures on a method in the theory of exponential sums. Tata Inst. of Fund. Res. Lectures on Math. and Physics, 80. Bombay, 1987.
- [14] I. Katai, The number of lattice points in a circle. (Russian) Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **8**(1965), 39–60.
- [15] T. Meurman, On the mean square of the Riemann zeta-function. Quart. J. Math. Oxford Ser. (2) 38 (1987), no. 151, 337–343.
- [16] W. Müller, On the asymptotic behaviour of the ideal counting function in quadratic number fields. Monatsh. Math. 108 (1989), no. 4, 301–323.
- [17] K. C. Tong, On divisor problem III, Acta math. Sinica **6** (1956), 515-541.
- [18] Kai-Man Tsang, Higher-power moments of $\Delta(x)$, $E(t)$ and $P(x)$, Proc. London Math. Soc.(3)**65**(1992), 65-84.
- [19] G. Voronoi, Sur une fonction transcendante et ses applications a la sommation de quelques séries, Ann. Sci. École Norm Sup. (3)**21**(1904), 207-267, 459-533.
- [20] Wenguang Zhai, On higher-power moments of $\Delta(x)$, Acta Arith. Vol.**112**(2004), 1-24.